

## ABSTRACT

A process for preparing (R)-5-[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]-5-hydroxy-3-oxo-1-heptanoic acid, tert-butylester comprising:

- (a) reduction of 5-[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]-3-oxo-1-pentanoic acid, (R)-2-hydroxy-1,2,2-triphenylethyl ester;
- 10 (b) hydrolysis of (R)-5-[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]-3-hydroxy-1-pentanoic acid, (R)-2-hydroxy-1,2,2-triphenylethyl ester using an alkali base in a solvent to form the acid;
- 15 (c) alkylation of the acid forming (R)-5-[2-(4-fluorophenyl)-5-(1-methylethyl)-3-phenyl-4-[(phenylamino)carbonyl]-1H-pyrrol-1-yl]-5-hydroxy-3-oxo-1-heptanoic acid, tert-butylester.